

Dissolved Nutrient, Chlorophyll-*a* and DOC Dynamic Under Distinct Riverine Discharges and Tidal Cycles Regimes at the Paraíba do Sul River Estuary, R.J., Brazil

G.C.T. Krüger†; C.E.V de Carvalho† and M. S. Suzuki†

Universidade Estadual do Norte Fluminense,
Centro de Biociências e Biotecnologia,
Laboratório de Ciências Ambientais,
Campos dos Goytacazes, 28013-602, R.J, Brasil.
carvalho@uenf.br



ABSTRACT

KRÜGER, G.C.T.; CARVALHO, C.E.V. and SUZUKI, M.S., 2006. Dissolved nutrient, Chlorophyll-*a* and DOC dynamic under distinct riverine discharges and tidal cycles regimes at the Paraíba do Sul river estuary, R.J., Brazil. Journal of Coastal Research, SI 39 (Proceedings of the 8th International Coastal Symposium), 724 - 730. Itajai, SC, Brazil, ISSN 0749-0208.

The aim of the present work is to evaluate the role of Paraíba do Sul River (PSR) in the nutrient, dissolved organic carbon and chlorophyll *a* input to the adjacent coastal area. In order to achieve this goal, four samplings were performed between 2000 and 2001, two in the dry season and two in the rainy season. Samples were collected every hour during a complete tidal cycle as well as physico-chemical parameters and currents direction and velocity with an acoustic Doppler profiler (ADP1500®). Inorganic (P-PO₄, N-NO₃, N-NO₂, N-NH₄ and Si) and organic (DON, DOP) nutrient as well as dissolved organic carbon (DOC) analysis was performed at the laboratory. Statistical difference between dry and rainy season were observed for the majority of the studied nutrients (Si, N-NO₃, N-NO₂, N-NH₄ and P-PO₄), COD and chlorophyll *a*. The average nutrient concentration observed at the present study were higher than other small and medium size rivers estuaries along the Brazilian south and southeastern coast, as the Camburiú (SC), except for NH₄ and PO₄, and the Mucuri (BA) and similar to the values observed for the Patos Lagoon estuary (RS).

ADDITIONAL INDEX WORDS: *Tropics, south America, coastal areas.*

INTRODUCTION

The high primary production and the position between the marine and the continental environment make the estuarine systems one of the most vulnerable ecosystems to the global changes and human activities (HASSEN, 2001).

Scientific studies in tropical estuaries began to be published at the literature between the sixties and the seventies and in the last years these areas are receiving special attention (BERNER and RAO, 1994; EYRE, 1995), mainly due to the riverine systems of these areas represent the main freshwater source to the world oceans (MILLIMAN and MEADE, 1983). The majority of the studies dealing with nutrient dynamic and flux to the coastal zones performed at the eastern South American Coast only considered the two large rivers of the region the Amazon and the São Francisco. Therefore there is a huge deficiency of data from the other medium and small sized rivers that also discharges their waters along this part of the Atlantic Ocean (CARNEIRO 1998; CARVALHO *et al.*, 2002).

Tropical estuaries present several differences from temperate and polar estuaries, the main are: the higher rainfall rates; the presence of perennial vegetation that reduces the soil erosion rates and a limited temperature variation. Besides that, the rivers from these regions present high water discharge; transport a smaller amount of suspended particulate matter (mainly due to the vegetation presents a higher density) and present a higher nutrient concentration (MEYBECK, 1984; NITTROUER *et al.*, 1995; ZHANG, 1999).

The world population growth has severely raised the demand on the aquatic and terrestrial systems. One third to half of the earth surface have already been changed due to the intense urbanization, agriculture, deforestation and reforestation processes that altered the hydrological cycle. Additional factors are the alteration of the natural biological communities and the introduction of exotic (non-native) species in the environments (VITOUSEK *et al.* 1997 apud SMITH *et al.*, 1999).

In Brazil the largest population densities were observed at coastal regions, being the PSR drainage basin entirely situated in an intense urbanized region. The main pollution sources to

the PSR lower basin include the disposal of untreated domestic effluents and the sewage from the sugarcane and alcohol industries also disposed directly at the river or in its affluent without any previous treatment (COSTA, 1998).

The present study objectives are to determinate the nutrient, chlorophyll *a* and DOC dynamic at the Paraíba do Sul River estuary during distinct riverine discharges (dry and rainy season) and tidal regime (spring and neap tides).

METHODS

The Paraíba do Sul River (Figure 1) is located between 20°26' and 23°38' S and 41° and 46°39' W, being entirely inserted at the southeastern region of Brazil. Its drainage basin is closely related to the most industrialized region of the country, the States of São Paulo, Minas Gerais and Rio de Janeiro.

The RPS river net discharge near its delta present a seasonal variation ranging from 300 to 1650 m³s⁻¹, with the winter months (June-August) corresponding to the lower discharge and the summer months (December-February) to the higher discharge period. The annual average discharge is 672 m³s⁻¹ (COSTA, 1998).

Four tidal cycles, one during neap tide and another during spring tide at October 2000 (dry season) and February 2001 (rainy season) were analysed. In each survey water samples were taken (surface and bottom), physico-chemical parameters were measured (pH, conductivity, temperature, dissolved oxygen) every hour. Water samples were collected by a "Van Dorn" sampler and immediately filtered in cellulose acetate membranes (0.45µm) for nutrient analysis, and in glass fiber filters (GF/FV Millipore®) for DOC and chlorophyll *a*. For nutrients the filtered samples were than stored in plastic flasks and frozen until analysis. For DOC analysis the filtered samples were stored in amber glass flasks with 10 ml of H₃PO₄ 10% and stored at a refrigerator until analysis (CALASANS *et al.*, 1995).

At the laboratory the DOC was analyzed by high temperature catalytic oxidation (680°C), in a carbon analyzer Shimadzu TOC-5000 (SUGIMURA and SUZUKI, 1988). The dissolved




Figure 1. Map showing the geography of the Paraíba do Sul lower basin and the sampling site location (pointer).

nutrient species and the chlorophyll *a* were analyzed by colorimetric methods. The ions N-NO_3 and N-NO_2 were analyzed by flow injection analysis (ASIA Ismatec) and the ions P-PO_4 , N-NH^+ and Si analyzed following the methodology described by GRASSHOFF et al. (1983). The DOP and the DON were obtained by the subtraction of the total concentration from the inorganic fraction. All the analysis was performed in triplicate.

In order to detect significant statistical differences between the surface and the bottom of the water column and between the dry and the rainy periods ($\alpha = 0,05$), a T test for independent samples was used with the help of the software Statistica for Windows, version 5.5, 2000 (by StatSoft, inc.)

RESULTS

Hydrodynamic

During the dry period samplings (October) the average salinity was 6.36, ranging between 0.1 and 29.5 psu, with higher values at the bottom samples. In the rainy period (February) on the neap tide sampling, there was no penetration of the coastal waters inside the estuary and thus no salinity was observed. Although during the spring tide the salinity ranged from 0.04 to 30.37 psu (Figure 2). Concerning the tide amplitude, the neap

tides presented smaller variations when compared with the studied spring tides. During the rainy season sampling the stream velocity ranged from 1.44 m s^{-1} to 1.28 m s^{-1} , although during the neap tide sampling all the measures velocities were negative indicating a seaward flow. In the dry period the variation was smaller, with stream velocity ranging from 0.76 to 0.78 m s^{-1} (Figure 3). The average pH values were 8.21 in the dry period and 7.22 in the rainy. The average dissolved oxygen was also higher during the dry period (6.38 mg l^{-1}) when compared to the rainy period (4.64 mg l^{-1}). The average temperature ranged from 23.8°C to 27.3°C in October and from 26.0°C to 31°C in February.

Nutrients and Chlorophyll *a*

Table 1 summarizes the obtained results during all the sampling periods. In general, a decrease in the nutrient concentration was always observed during the flood and at the high tide. The NO_3 correspond to 94% to 95% of the dissolved inorganic nitrogen (DIN). In the dry period the dissolved organic nitrogen (DON) was the dominant fraction, although in the rainy season the NO_3 concentrations were slightly higher. Both DON and NO_3 values decreased their concentrations with the salinity increase. The ammonia concentrations were generally low in the estuary, and in the dry period (October), and




Figure 2. Salinity variations along time during neap and spring tides in the rainy and dry season.

during the high tide the concentrations were below the detection limit of the method. The NO_2 and the NH_4 presented an inverse trend when compared with NO_3 and DON, with situations of gain in relation to the salinity (Figure 4). The orthophosphate concentrations were always higher than the dissolved organic phosphate (DOP) in all the studied periods and tides, both elements presented higher concentrations during the ebb tide. In relation to Si, the rainy period presented higher concentrations when compared to the dry period, probably reflecting the fluvial input of this element to the estuarine system. The DOC variation was better understood during the spring tides where the concentrations tend to raise at the same time that the tide was ebbing. The DOC concentrations ranged from 1.5 mg l^{-1} to 5.6 mg l^{-1} . Orthophosphate presented the same trend observed for silica and nitrogen, higher concentrations observed in the rainy period with the predominance of the inorganic form.

The N:P ratio ranged from 16 to 39, with a relative




Figure 3. Average stream velocity along time during neap and spring tides in the rainy and dry season.

enrichment of the inorganic nitrogen in relation to the orthophosphate.

Chlorophyll *a* presented its higher values when the river discharge was lower (dry period). During the dry period the chlorophyll *a* decrease its concentration with the penetration of the marine waters inside the estuary (flood tide). Although during the rainy season an opposite trend was observed, with increasing concentrations while the tide was flooding.

Statistical Analysis

The T-test showed significant statistical differences ($P < 0,05$) between the surface and bottom samples for temperature in both studied periods (dry and rainy), with higher values observed at the surface ($\pm 1^\circ\text{C}$). Concerning the nutrients, only NO_3 presented significant statistical differences ($P < 0,02$) between surface and bottom samples on the February sampling (rainy season, neap tide). Comparing the nutrient, Chl-*a* and DOC

Table 1. Average nutrient, chlorophyll *a* and DOC concentrations in the Paraíba do Sul River Estuary, in the rainy and dry seasons.

	COD (mg l^{-1})	Chl- <i>a</i> ($\mu\text{g l}^{-1}$)	P- PO_4 (μM)	POD (μM)	NO_3 (μM)	NO_2 (μM)	NH_4 (μM)	NOD (μM)	Si(OH) $_4$ (μM)	N:P
<i>19 October 2000 (dry season, neap tide) (n=28)</i>										
Mean	3,24	15,91	0,66	0,25	20,7	0,36	0,50	-	132,2	35
s.d.	0,4	4,26	0,18	0,17	6,2	0,09	0,43	-	26,8	17
Range	1,9-4,6	7,1-33,7	0,25-1,15	0,08-0,63	4,2-30,4	0,18-0,64	0,05-1,27	-	30,7-214,7	19-7
<i>25 October 2000 (dry season, spring tide) (n=28)</i>										
Mean	3,01	19,91	0,65	0,54	7,41	0,57	0,33	21,76	67,4	16
s.d.	0,6,5	7,19	0,31	0,15	5,64	0,30	0,39	6,23	33,2	14
Range	1,2-4,6	3,0-37,3	0,25-2,03	0,27-1,15	0,46-15,60	0,24-1,14	0,05-1,33	40,7-6,55	14,3-103,0	3-41
<i>17 February 2001 (rainy season, neap tide) (n=28)</i>										
Mean	3,19	2,91	1,19	0,57	32,8	0,20	1,25	11,85	174,0	39
s.d.	0,79	0,98	0,3	0,51	4,2	0,19	0,74	7,48	26,38	24
Range	2,3-6,7	0,92-5,55	0,56-1,98	0,1-2,9	24,2-37,8	0,10-0,84	0,31-3,94	2,1-44,9	62,6-205,5	20-89
<i>24 February 2001 (rainy season, spring tide) (n=28)</i>										
Mean	2,19	8,41	0,69	0,35	17,56	0,32	0,58	19,90	123,6	29
s.d.	0,38	2,58	0,29	0,29	11,49	0,22	0,38	16,23	53,15	15
Range	1,4-3,7	1,9-23,4	0,16-1,48	0,04-1,02	2,71-33,75	0,08-0,86	0,08-2,45	1,8-57,4	21,2-203,9	5-46

s.d.: standard deviation, n: number of samples.




Figure 4. NH_4 and NO_2 variation during the spring tides (25/10/00).

behavior between the two studied periods (dry x rainy, Table 2), the test was statistically significant for all the studied elements; the only exception was the DOP ($P=0,13$) and the DON ($P=0,10$). Concerning the physico-chemical parameters, it was observed seasonal variation for pH and temperature. The average pH was 8.21 and 7.22 in the dry and rainy periods respectively, and the average temperature was 26.3°C in October/2000 and 29.5°C in February/2001. All the data was statistically correlated using the Pearson coefficient ($P<0.05$). Based on this data it was possible to show that in the dry period the penetration of the marine waters inside the estuary is probably promoting the decrease of PO_4 (-0,44), NO_3 (-0,74) and Si (-0,75). During the rainy period the salinity presented a significant correlation with NO_3 (-0,62), Si (-0,71) and with DOC (-0,54).

DISCUSSION

The hydrographic conditions of the Paraíba do Sul River estuary are closely related with the seasonal variations. As well as in other Brazilian and world rivers and bays, it's accepted that nutrient concentrations have risen over the last 50 years as a consequence of agricultural practices (excess of fertilizers) and the discharges of urban wastewaters (SIERRA *et al.*, 2002; VALIELA and BOWEN, 2002). As already observed in other tropical estuaries, there is a general trend showing an increase in the nutrient concentration in the rainy season mainly due to the riverine input that also raises during this time of the year (EYRE, 1994). Once mixed with marine waters, inorganic nutrients are progressively transformed along the salinity gradient and, the complex biogeochemical processes occur in the low salinity (0-10) (RAGUENEAU *et al.*, 2002). During periods of high riverine discharges estuarine water column is expected to be highly stratified, this trend was not observed in the Paraíba do Sul Estuary. The main reasons for the water column homogeneity are probably the low depths of the estuary (maximum depth of the estuary channel is 6.5m) and the adjacent continental shelf; The low tidal amplitude (spring tide average variation is 0.65 m) and the presence of sand banks in front of the estuary mouth. Therefore, according to SCHETTINI (2001) although dominated




Figure 5. N:P ratio distribution during the dry season spring tide.

Table 2. *P* values obtained with the T-test for nutrients, DOC and Chlorophyll *a*. *P* 0,05 in bold represent significant correlation at 5% confidence level between dry and rainy periods.

T- Test	
DOC	0,006
P- PO_4	0,0000
DOP	0,130
N- NO_3	0,00003
N- NO_2	0,00033
DON	0,108
N- NH_4	0,0005
Si (OH) ₄	0,0000
Chl-a	0,0000

by fluvial waters the friction forces on the PSR estuary are sufficient to maintain the homogeneity of the water column practically all the time.

Nutrient Dynamic in Dry Season

According to several authors (SILVA, 2000; TIAN *et al.*, 1993; KRESS *et al.*, 2002) during the dry period the primary productivity tend to be enhanced mainly due to a lower turbidity of the system, causing a raise of the Chl-*a* and dissolved oxygen concentrations in the water column.

During this period the DON was predominant nitrogen from in the PSR estuary, probably reflecting the raise in the activities of sugarcane and alcohol factories, enhancing the effluent discharges with high organic matter content to the riverbed. In addition the lower water level decreases the dilution effect (FIGUEIREDO, 1999). The interference of the salt wedge on the DON concentrations were better observed during the spring tide (25/10/00) when the values drop from 50.65 μM to 16.41 μM three hours after the beginning of the ebb tide. The dilution effect observed for DON is indicating a conservative behavior, the same pattern was observed for NO_3 , which decreased its concentrations from 16.84 μM when the salinity was 2.0 psu to 0.46 μM when the salinity reaches 20.34 psu. CARNEIRO (1998) also observed the same behavior in the Paraíba do Sul Estuary in March and July 1995. PAKULSKI *et al.* (2000), based in the DIN and DON distribution, suggested that the increment of DON in relation to the DIN in intermediate salinities is probably associated with the transformation of the nitrate in DON mediated by the phytoplankton assimilation. The average NO_3 concentrations obtained in this study were higher than the values observed for other tropical estuaries during the dry period as the Moresby River in Australia (EYRE, 1994) and the Nicoya Gulf in Costa Rica (KRESS *et al.*, 2002), both under environmental stress. Concerning the NH_4 distribution the low observed concentrations followed the same pattern described for the Nicoya Gulf, which presented negligible values (KRESS *et al.*, 2002). A tendency of increasing concentrations following the salinity gradient was observed for NO_2 and NH_4 (Figure 5), is probably due to the release of interstitial waters from the estuarine sediments that are enriched in these elements (Turner, 1990) was also observed by CARNEIRO (1998) in the Paraíba do Sul Estuary.

Phosphate concentrations presented a slightly increase in low salinity (2 - 5 psu) decreasing its concentrations in higher salinities. The mechanisms that include the PO_4 dynamics and its other chemical forms in the Paraíba do Sul estuary seems to be more related to chemical processes, considering the salinity gradient and the pH changes. Many authors suggested that a buffer mechanism is probably regulating the PO_4 concentrations between 0.6 - 1.4 μM in estuarine systems. This mechanism is based in the adsorption/desorption of the PO_4 from the suspended particulate matter and sediments when the concentrations are low. The process is reversible and mainly controlled by the pH variation (below 7.8 the PO_4 removal is intensified), followed by other factors as the SPM and

sediments concentration, biological activity and residence time (EYRE, 1994; FOX *et al.*, 1986; FANG, 2000; RAGUENEAU *et al.*, 2002). Concerning the PO_4 adsorption on iron hydroxides, assumed to be the most important inorganic phosphorous removal process between salinities of 5–15 psu (FOX, 1990; FANG, 2000) the absence of the iron concentrations data do not allowed any speculations about it. Although the presence of iron enriched soils in the PSR drainage basin may contribute to the iron input for the estuary, probably becoming an important factor in the phosphate removal process (CARVALHO *et al.*, 2002). The dissolved organic fraction (DOP) is probably associated with the raise of the organic matter during this period of the year, as indicated by the COD distribution. The other possible DOP source to the estuary is probably the primary production, also higher during the dry period (FANG, 2000; CARNEIRO, 1998). Finally the increasing DIP concentration in the water column is probably reflecting the sediment resuspension and the orthophosphate desorption, once the DOP values are low and a possible conversion of DOP into DIP seems to be improbable due to the small importance of the organic fraction in the total dissolved phosphorous. The phosphorous values measured in the present study are in the same range of other studies developed in tropical estuarine systems of Brazil and other tropical countries (FANG, 2000; KRESS *et al.*, 2002), STERZA, 2002; PEREIRA-FILHO *et al.*, 2001 and SIERRA *et al.*, 2002).

The COD that generally present a conservative behavior in the PSR estuary present a clear tendency to raise its concentrations with the ebb tide (3,15 mg.l^{-1} when the salinity was 5,8 psu increasing to 4,0 mg.l^{-1} when the salinity reaches 1,7 psu). The COD behavior, as well as the concentration range follows other studies previously developed in the PSR Estuary (CALASANS *et al.*, 1995 and CARNEIRO, 1998)

Although high Chl-a concentrations were measured, the Si concentrations were lower than the observed during the rainy period, probably indicating that the main Si source for the estuary are the intemperism processes, as soil lixiviation and the flood of marginal inundated soils, that generally occurs during high river discharge periods (KRESS *et al.*, 2002; SILVA, 2000).

Nutrient Dynamic in Rainy Season

With the beginning of the rainy season and the raise of the river discharge, the river became the most important source of nutrients for the estuary. During the rainy season the mixing of the distinct water masses play a secondary role as source of the studied nutrient, mainly due to the low tidal amplitude and the homogeneity of the water column.

The Chl-a distribution followed a distinct pattern when compared with the dry period. Peaks of high Chl-a concentrations were observed just after the peak of the flood tide. This behavior have already been observed by PEREIRA-FILHO *et al.* (2001) for the Camburiú River Estuary (SC, Brazil), and the probable explanation for this behavior is the occurrence of a very high productive zone near the estuarine mouth. During the flood tide part of this phytoplankton enriched water is probably entering the main estuarine channel causing the increment of the Chl-a after the flood tide.

The NO_3 became the main nitrogen form present in the estuary, been basically related to the input from the surface runoff for the riverine system. According to YIN *et al.* (1995) and HERNANDEZ-AYON *et al.* (1993), that studied estuaries in Canada and in the USA, when the high river discharge period started no physical or biological processes (e.g. mineralization and the phytoplankton activity) are capable to alter the NO_3 concentration in estuarine waters. According to CARNEIRO (1998) the NO_3 in the PSR estuary is quickly transported trough the estuary not being available to the reactions and transformations along the mixing zone. The same situation could probably be applied for DON and NH_4 that most of the time presented a conservative behavior, with moments of gain and lost without a clear trend. In addition, the low Chl-a concentrations did not support the hypothesis of a significant

release of this nitrogen forms from primary producers, to the water column. Concerning the NO_2 and NH_4 distribution, peaks of these elements were observed in the end of the flood tide, and are directly correlated with the decrease in the dissolved oxygen concentrations, probably suggesting the reduction of the NO_3 associated with the bottom sediments (ZHANG *et al.*, 1999).

Higher PO_4 concentrations are generally expected in the rainy season mainly due to the input caused by the surface runoff along the drainage basin as well as by the flood of the plain river margins, where cattle farming and agriculture are developed, enhancing the transport of the material accumulated during the dry season to the river bed (SILVA, 2000). In the rainy season a significant correlation between PO_4 and pH was observed (0.43), this correlation is probably indicating that the PO_4 removal mediated by the pH that was described for the dry period is also occurring. The dissolved organic form (DOP) presented peaks of concentration during the flood tide, this fact in addition to the Chl-a data is probably indicating that a small fraction of the fluvial PO_4 is being transformed in DOP by the assimilation and transformation caused by the primary producers (PAKULSKI *et al.*, 2000).

Although DOC concentration were expected to be higher during the rainy season (CARNEIRO, 1998; FIGUEIREDO, 1999), in the present study the values were a little bit higher during the dry season. This trend was probably due to the more intense penetration of the salt wedge in 24/02 (spring tide), when the DOC concentrations decreased during the flood tide, from 2.5 mg.L^{-1} to 1.5 mg.L^{-1} . Besides that, the PSR discharge during this sampling was lower than the historical average for the rainy season. Therefore a statistical difference was expected between dry and rainy season, although not with lower concentrations during the rainy season.

BESSA and PAREDES (1990) studying the São Francisco River already observed lower DOC concentrations during the dry season. These authors attributed this behavior to the intense evaporation of the river waters, and consequently increase of the organic matter, as well as the decomposition of the resuspended particulate organic matter.

The silica values were inside the range already observed for some estuarine systems (CABEÇADAS *et al.*, 1999; TIAN *et al.*, 1993) and higher than other estuaries (HASSEN, 2001; HERNANDEZ-AYON *et al.*, 1993; PAKULSKI *et al.*, 2000; PEREIRA-FILHO *et al.*, 2001 and CORBETT *et al.*, 2002). Although Si presented a conservative behavior, a significant correlation with NO_3 (0.61) is probably indicating that part of the silica concentration is being removed by the biological assimilation (EYRE, 1994).

Concerning the tidal influence, Si concentrations tend to decrease with increasing salinity (CORBETT *et al.*, 2002; PEREIRA-FILHO *et al.*, 2001; RAGUENEAU *et al.*, 2002).

Finally, according to Zhang (1999), the Si/N ratio that could interfere in the primary production ($= 0,5 - 1$), variate between 5 and 7 during all samplings, suggesting that silica is not a limiting nutrient for the system. According to previous studies, the N:P ratio for the Paraíba do Sul river (62 to 496) was much higher than the PSR estuary where it variates between 9 and 23 (CARNEIRO, 1998). In the estuarine zone under the influence of the mixing waters the observed ratio was lower (Figure 5), mainly due to the strong phosphorous desorption when compared with the DIN. The values observed in this study are inside of the range reported for others estuaries like the Vitória Bay (ES), 10–21 (STERZA, 2002), Camburiú River Estuary (SC) 12–149 (PEREIRA-FILHO *et al.*, 2001) and the Danubio River Estuary, 40–100 (RAGUENEAU *et al.*, 2002), all of them with relative anthropogenic influence (domestic sewage, fertilizers, etc.).

CONCLUSIONS

The elements concentrations showed that the rain events are the main regulators of the nutrient, Chl-a and DOC dynamic in the PSR estuary. With exception of the PO_4 , NO_2 and NH_4 the majority of the studied elements presented a conservative

behavior, being the decrease of its concentrations attributed to the dilution effect caused by the mixture of marine and fluvial waters. The presented data showed that the freshwater end-member composition are comparable with other Brazilian and world rivers. The nutrient ratios could be considered as low for a polluted river, and they did not seem to increase the estuarine primary production. Independently from the river discharge, during the spring tides higher amplitudes between maximum and minimum nutrient concentrations were observed (with exception of DOC and NH₄). An influence of the tides in the primary productivity was also observed, with higher variation of the Chl-a concentrations also observed during the spring tides.

In the period of lower fluvial discharge the estuary presented the highest concentrations of Chl-a and the organic nutrient forms. When the discharge increased the river became the main source of nutrients for the estuary. In this period the photosynthetic activity also decreased and a consequently decrease in the dissolved oxygen was also observed. Finally, during the rainy period the decrease in the DON concentrations are probably associated with the mineralization processes indicated by the increase of the NH₄ concentrations during this period.

ACKNOWLEDGEMENTS

This work was supported by FAPERJ (Proc. N° E26-170.368/199) and CNPq (TMTM Proc. N° 103415/00-9). The authors also would like to thank M.Sc. Wanderson Primo de Souza, Dr. Alexandre Gomes Ferreira, M.Sc. Glaucete Magalhães Gonçalves and B.Sc. André Luis dos Santos Machado for helping during the samplings.

LITERATURE CITED

- BERNER, R.A. and RAO, J.L., 1994. Phosphorus in sediments of the Amazon river and estuary: implications for the global flux of phosphorus to the sea. *Geochimica et Cosmochimica Acta*, 58, 2333-2340.
- BESSA, M.F. and PAREDES, J.F., 1990. Transporte de carbono e do nitrogênio orgânico e inorgânico dissolvidos pelo rio São Francisco, Brasil, Durante um ano hidrológico (1984-1985). *Geochimica Brasiliensis*, 4(1): 17-31.
- CALASANS, C.F.; CARVALHO, C.E.V.; OVALLE, A.R.C.; CORREIA, V.G., and REZENDE, C.E., 1995. Distribuição de carbono orgânico dissolvido na porção inferior do Rio Paraíba do Sul, Rio de Janeiro. V Congresso Brasileiro de Geoquímica e III Congresso de Geoquímica dos países de língua portuguesa (Niterói, Rio de Janeiro).
- CARNEIRO, M.E.R., 1998. Origem, transporte e destino da matéria orgânica no estuário do Rio Paraíba do Sul. Niterói, Rio de Janeiro: Universidade Federal Fluminense, Ph.D. thesis, 210p.
- CARVALHO, C.E.V.; SALOMAO, M.S.M.B.; MOLISANI, M.M.; REZENDE, C.E., and LACERDA, L.D., 2002. Contribution of a Medium-Sized Tropical River to the Particulate Heavy Metal Load for the South Atlantic Ocean. *The Science of the Total Environment*, 284(1-3): 85-93.
- CORBETT, D.R.; DILLON, K.; BURNETT, W., and SCHAFER, G., 2002. The spatial variability of nitrogen and phosphorus concentration in a sand aquifer influenced by onsite sewage treatment and disposal systems: a case study on St. George Island, Florida. *Environmental Pollution*, 117, 337-345.
- COSTA, H., 1998. Uma avaliação da qualidade das águas costeiras do Estado do Rio de Janeiro. Rio de Janeiro: Fundação de Estudos do Mar, 261p.
- EYRE, B., 1994. Nutrient biogeochemistry in the tropical Moresby river estuary system north Queensland, Australia. *Estuarine, Coastal and Shelf Science*, 39, 15-31.
- EYRE, B.D., 1995. A first-order nutrient budget for the tropical Moresby estuary and catchment, North Queensland, Australia. *Journal of Coastal Research*, 11, 717-732.
- FANG, T.H., 2000. Partitioning and behavior of different forms of phosphorus in the Tanshui estuary and one of its tributaries, northern Taiwan. *Estuarine, Coastal and Shelf Science*, 50, 689-701.
- FIGUEIREDO, R.O., 1999. Transporte de Carbono e Nitrogênio no Baixo Paraíba do Sul: Fluxos e Processos. Campos dos Goytacazes, Rio de Janeiro: Universidade Estadual do Norte Fluminense, Ph.D. thesis, 177p.
- FOX, L.E.; SAGER, S.L., and WOFSEY, S.C., 1986. The chemical control of soluble phosphorus in the Amazon estuary. *Geochimica et Cosmochimica Acta*, 50, 783-794.
- GRASSHOF, K.; EHNHARDT, M. and KREMILING, K., 1983. *Methods of Seawater Analysis*. Weinheim: Verlag Chemie, 419p.
- HASSEN, M.B., 2001. Spatial and temporal variability in nutrients and suspended material processing in the Fier d'Ar Bay. *Estuarine, Coastal and Shelf Science*, 52, 457-469.
- HERNANDEZ-AYON, J.M.; GALINDO-BECT, M.S.; FLORES-BAEZ, B.P., and ALVAREZ, S., 1993. Nutrient concentration are high in the turbid waters of Colorado river delta. *Estuarine, Coastal and Shelf Science*, 37, 593-602.
- KRESS, N.; COTO, S. L.; BRENES, C. L.; BRENNER, S., and ARROYO, G., 2002. Horizontal transport and seasonal distribution of nutrients, dissolved oxygen and chlorophyll-a in the Gulf of Nicoya, Costa Rica: a tropical estuary. *Continental Shelf Research*, 22, 51-66.
- MEYBECK, M., 1982. Carbon, nitrogen and phosphorus transport by major world rivers. *American Journal Science*, 282, 401-501.
- MILLIMAN, J.P. and MEADE, P.H., 1983. World-wide delivery of riverine sediment to the ocean. *Journal of Geology*, 91, 1-21.
- NITTROUER, C.A.; BRUNSKILL, G.J., and FIGUEIREDO, A.G., 1995. Importance of tropical coastal environment. *Geomarine Letters*, 15, 121-126.
- PAKULSKI, J.D.; BENNER, R.; WHITLEDGE, T.; AMON, R.; EADIE, B.; CIFUENTES, L.; AMMERMAN, J., and STOCKWELL, D., 2000. Microbial metabolism and nutrient cycling in the Mississippi and Atchafalaya plumes. *Estuarine, Coastal and Shelf Science*, 50, 173-184.
- PEREIRA-FILHO, J.; SCHETTINI, C.A.F.; RÖRIG, L., and SIEGLE, E., 2001. Intratidal variation and net transport of dissolved inorganic nutrients, POC and chlorophyll a in the Camboriú river estuary, Brazil. *Estuarine, Coastal and Shelf Science*, 53, 249-257.
- RAGUENEAU, O.; LANCELOT, C.; EGOROV, V.; VERVLIMMEREN, J.; COSIASU, A.; DÉLIAT, G.; KRASSTEV, A.; DAOUD, N.; ROUSSEAU, V.; POPOVITCHEV, V.; POPA, L., and CAUWET, G., 2002. Biogeochemical transformations of inorganic nutrients in the mixing zone between the Danube river and the north-western Black sea. *Estuarine, Coastal and Shelf Science*, 54, 321-336.
- SCHETTINI, C.A.F.; TRUCCOLO, E.C., and CARVALHO, C.E.V., 2002. Caracterização hidrodinâmica da passagem sul do estuário do rio Paraíba do Sul. I Congresso Brasileiro de Oceanografia (Rio de Janeiro).
- SIERRA, J.P.; SÁNCHEZ-ARCILLA, A.; GONZÁLES DEL RIO, J.P.; FLOS, J.P.; MUVELLÁN, E.; MÖSSO, C.; MARTÍNEZ, R.; RODILLA, M., and FALCO, S.P., 2001. Spatial distribution of nutrient in the Elbro estuary and plume. *Continental Shelf Research*, 22, 361-378.
- SILVA, C.F., 2000. Hidrogeoquímica fluvial da porção inferior do rio Paraíba do Sul, R.J. Campos dos Goytacazes, Rio de Janeiro: Universidade estadual do Norte Fluminense, Master's thesis, 65 p.
- SMITH, V.H.; TILMAN, G.D., and NECOLA, J.C., 1999. Eutrophication impacts of excess nutrients inputs on freshwater, marine and terrestrial ecosystems. *Environmental Pollution*, 100, 179-196.
- STERZA, J.M., 2002. Composição, abundância e distribuição espaço-temporal do zooplâncton relacionadas com a hidroquímica no sistema estuarino Baía de Vitória/Canal da Passagem, Vitória, ES. Campos dos Goytacazes, Rio de Janeiro: Universidade Estadual do Norte Fluminense, Master's thesis, 88 p.
- SUGIMURA, Y. and SUZUKI, Y. 1998. A high-temperature

- catalytic oxidation method for the determination of nonvolatile dissolved organic carbon in seawater by direct injection of a liquid sample. *Marine Chemistry*, 24, 105-131.
- TIAN, R. C.; HU, F. X., and MARTIN, J. M., 1993. Summer nutrients fronts in the Changjiang (Yantze river) estuary. *Estuarine, Coastal and Shelf Science*, 37, 27-4.
- VALIELA, I. and BOWEN, J.L., 2002. Nitrogen sources to watersheds and estuaries: role of land cover mosaics and losses within watersheds. *Environmental Pollution*, 118, 239-248.
- ZHANG, J.; YU, Z.G.; WANG, J.T.; REN, J.L.; CHEN, H.T.; XIONG, H.; DONG, L.X., and XU, W.Y., 1999. *Estuarine, Coastal and Shelf Science*, 49, 385-400.